



APHID ALERT SUMMARY

This alert summarises up-to-date results from the Rothamsted/SASA suction-trap (ST) network and the FERA yellow water-pan trap (YWT) network. Further details of the ST results can be found below and further details of the YWT results can be found at www.potato.org.uk/online-toolbox/aphid-monitoring.

GENERAL

Aphid flight activity dropped for a second consecutive week due to continued unsettled weather and rapidly mounting pressure from natural enemies.

CEREALS

Grain aphid (*Sitobion avenae*) numbers have increased a little this week as they leave ripening crops, but cereal aphid numbers overall remain low for the time of year. There have been no reports of significant numbers of cereal aphids in crops. The threshold for control is 66% of tillers infested from GS61 to two weeks before the end of grain filling, but many crops are clearly already beyond this stage. The first barley crops have been harvested in East Anglia and southern England on lighter land, but the winter wheat harvest is yet to begin.

POTATOES

Virus pressure continued to drop throughout the country, but remains high in the midlands and the south west, but low across Scotland. Numbers of the peach-potato aphid (*Myzus persicae*) have fallen again this week, and are now low everywhere. Numbers of the black bean aphid (*Aphis fabae*) are dropping, but this species continues to drive much of the virus pressure. This species is a good vector of PVA, so PVA-susceptible varieties may be especially at risk. Cereal aphids remain at low levels for the time of year. Further regional information on potato virus vectors can be accessed here: www.potato.org.uk/online-toolbox/aphid-monitoring.

OILSEED RAPE

Aphids are no longer an issue in most crops with harvesting well underway in East Anglia and southern England on lighter land.

FIELD BRASSICAS and LEAFY VEGETABLES

Numbers of the mealy cabbage aphid (*Brevicoryne brassicae*) have risen this week in central England. The currant-lettuce aphid (*Nasonovia ribisnigri*) was caught in the suction-traps at Writtle and Hereford.


CARROTS

The numbers of the willow-carrot aphid (*Cavariella aegopodii*) flying are now very low everywhere.

PEAS and BEANS

Field reports suggest numbers of the black bean aphid (*Aphis fabae*) remains high in crops across much of England. The threshold for spraying is 10% of plants infested where the pods are still filling. Numbers of the pea aphid (*Acyrtosiphon pisum*) have risen a little in our traps and reports indicate they are common in peas, particularly in eastern England. Combining peas should be sprayed when around 20% of plants are infested and vining peas when 15% are infested.

SUCTION-TRAPPING RESULTS



The information below relates to suction-trap samples collected during bulletin week 15:13/7 – 19/7.

- Grain aphid (*Sitobion avenae*) numbers have increased a little this week as they leave ripening crops, but cereal aphid numbers overall remain low for the time of year.
- Numbers of the peach–potato aphid (*Myzus persicae*) have fallen again this week, and are now low everywhere.
- The potato aphid (*Macrosiphum euphorbiae*) was caught in low numbers at eight sites.
- The numbers of the willow–carrot aphid (*Cavariella aegopodii*) are now low everywhere.
- The mealy cabbage aphid (*Brevicoryne brassicae*) was caught at ten sites this week, with high numbers at Wellesbourne (855) and Hereford (413).
- Numbers of the pea aphid (*Acyrtosiphon pisum*) increased at nine sites this week, with accumulated numbers highest at Starcross.
- Numbers of the black bean aphid (*Aphis fabae*) remain high, but are now decreasing across much of England.
- Numbers of aphid predators (hoverflies, ladybirds and lacewings) are increasing in our trap samples and field reports suggest are now much more common in the field.

Suction-trap sites

The tables below show current and accumulated totals with comparisons to previous years. ‘/’ indicates that identifications have not been completed and ‘*’ indicates where totals have been corrected proportionally to seven days, because fewer days’ samples having been identified.

Rose–grain aphid (*Metopolophium dirhodum*)

Rose–grain aphid (<i>Metopolophium dirhodum</i>)	Bulletin Week Totals 13/07-19/07				Accumulated until 19/07		
	2015	Compared to last Bulletin week	2014	10-year average 2005-14	2015	2014	10-year average 2005-14
Dundee	25*	↑	640	977	45	/	1631
Gogarbank (Edinburgh)	45	↑	24	139	79	169	499
Newcastle	0*	↓	30	59	16	/	149
York	5*	↓	N/A	N/A	49	N/A	N/A
Preston	2*	↓	8	26	75	/	355
Kirton	11	↑	/	134	163	/	1235
Broom’s Barn (nr Bury St Edmunds)	8	↑	1	102	231	133	1149
Wellesbourne	0*	↓	12	49	276	/	402
Hereford	21*	↓	10	79	272	91	489
Rothamsted (Harpenden)	4	↓	10	59	150	146	629
Writtle	0*	↓	5	64	103	236	639
Silwood Park (nr Ascot)	5*	↓	/	6	94	/	176
Wye	26*	↑	/	44	74	/	255
Starcross (nr Exeter)	4*	↓	21	13	392	933	342

The rose–grain aphid was caught at eleven sites this week, with a minor hotspot at Edinburgh (45), but numbers decreased at nine sites. Accumulated numbers are below the ten year average everywhere except Starcross.

Bird cherry–oat aphid (*Rhopalosiphum padi*)

Bird cherry–oat aphid (<i>Rhopalosiphum padi</i>)	Bulletin Week Totals 13/07-19/07				Accumulated until 19/07		
	2015	Compared to last Bulletin week	2014	10-year average 2005-14	2015	2014	10-year average 2005-14
Dundee	0*	↓	447	303	12	/	846
Gogarbank (Edinburgh)	0		101	65	5	336	720
Newcastle	0*		47	33	4	/	303
York	2*	↓	N/A	N/A	22	N/A	N/A
Preston	0*	↓	28	30	15	/	140
Kirton	16	↑	/	71	85	/	765
Broom's Barn (nr Bury St Edmunds)	5	↑	5	76	52	90	817
Wellesbourne	1*	↓	33	93	17	/	264
Hereford	5*	↑	41	53	31	118	251
Rothamsted (Harpenden)	3	↑	42	54	25	121	385
Writtle	6*	↑	22	52	75	178	472
Silwood Park (nr Ascot)	5*	↑	/	28	22	/	240
Wye	8*	↑	/	55	47	/	285
Starcross (nr Exeter)	2*	↓	68	72	78	292	408

The bird cherry–oat aphid was caught at ten sites in very low numbers. The accumulated numbers are below average.

Grain aphid (*Sitobion avenae*)

Grain aphid (<i>Sitobion avenae</i>)	Bulletin Week Totals 13/07-19/07				Accumulated until 19/07		
	2015	Compared to last Bulletin week	2014	10-year average 2005-14	2015	2014	10-year average 2005-14
Dundee	4*	↓	587	150	26	/	305
Gogarbank (Edinburgh)	18	↑	139	55	27	347	232
Newcastle	6*	↑	54	46	10	/	114
York	49*	↓	N/A	N/A	252	N/A	N/A
Preston	16*	↑	63	63	60	/	281
Kirton	4	↑	/	269	38	/	1092
Broom's Barn (nr Bury St Edmunds)	12	↑	5	220	60	149	986
Wellesbourne	22*	↓	78	268	88	/	826
Hereford	88*	↑	81	163	229	217	568
Rothamsted (Harpenden)	13	↑	99	315	58	309	1141
Writtle	14*	↑	92	270	66	553	1101
Silwood Park (nr Ascot)	16*	↑	/	100	75	/	542
Wye	8*	↑	/	450	36	/	685
Starcross (nr Exeter)	140*	↑	105	71	368	343	441

Numbers of grain aphid increased at thirteen sites this week, with hotspots at Hereford (88) and Starcross (140). The accumulated numbers are below average for the time of year.

Peach–potato aphid (*Myzus persicae*)

Peach–potato aphid (<i>Myzus persicae</i>)	Bulletin Week Totals				Accumulated until		
	2015	Compared to last Bulletin week	2014	10-year average 2005-14	2015	2014	10-year average 2005-14
Dundee	0*	↓	5	3	10	/	10
Gogarbank (Edinburgh)	3	↑	2	1	30	6	10
Newcastle	8*	↓	5	2	32	/	15
York	21*	↓	N/A	N/A	432	N/A	N/A
Preston	9*	↓	5	2	154	/	110
Kirton	8	↓	/	26	2330	/	412
Broom's Barn (nr Bury St Edmunds)	2	↓	1	49	2170	1351	728
Wellesbourne	15*	↓	5	6	1194	/	368
Hereford	7*	↓	3	10	697	105	121
Rothamsted (Harpenden)	39	↑	3	74	1051	520	427
Writtle	4*	↓	5	53	1661	1158	751
Silwood Park (nr Ascot)	7*	↑	/	2	129	/	56
Wye	9*	↓	/	16	588	/	190
Starcross (nr Exeter)	11*		4	7	891	168	90

The peach–potato aphid was caught at thirteen sites this week. Numbers fell at ten sites, but the accumulated totals are above the 10 yr averages throughout the Country.

Potato aphid (*Macrosiphum euphorbiae*)

Potato aphid (<i>Macrosiphum euphorbiae</i>)	Bulletin Week Totals				Accumulated until		
	2015	Compared to last Bulletin week	2014	10-year average 2005-14	2015	2014	10-year average 2005-14
Dundee	2*	↓	5	6	8	/	20
Gogarbank (Edinburgh)	2	↑	2	4	13	27	47
Newcastle	1*	↑	2	4	12	/	19
York	0*	↓	N/A	N/A	19	N/A	N/A
Preston	0*		0	1	40	/	28
Kirton	0	↓	/	6	75	/	65
Broom's Barn (nr Bury St Edmunds)	0	↓	0	4	19	29	48
Wellesbourne	6*	↑	0	5	50	/	42
Hereford	12*	↑	0	9	98	31	78
Rothamsted (Harpenden)	6	↑	1	3	23	13	26
Writtle	3*	↑	0	3	51	35	46
Silwood Park (nr Ascot)	1*	↑	/	2	29	/	18
Wye	0*		/	2	6	/	19
Starcross (nr Exeter)	0*	↓	1	1	56	21	40

The potato aphid was caught at eight sites this week in low numbers, with highest numbers at Hereford (12).

Mealy Cabbage aphid (*Brevicoryne brassicae*)

Cabbage aphid (<i>Brevicoryne brassicae</i>)	Bulletin Week Totals 13/07-19/07				Accumulated until 19/07		
	2015	Compared to last Bulletin week	2014	10-year average 2005-14	2015	2014	10-year average 2005-14
Dundee	0*		0	1	0	/	3
Gogarbank (Edinburgh)	0		0	0	0	0	3
Newcastle	0*		0	0	4	/	8
York	14*		N/A	N/A	109	N/A	N/A
Preston	0*	↓	2	1	7	/	25
Kirton	50	↓	/	12	920	/	93
Broom's Barn (nr Bury St Edmunds)	10	↓	0	29	188	43	260
Wellesbourne	855*	↑	9	17	4682	/	376
Hereford	413*	↑	4	8	1191	11	142
Rothamsted (Harpenden)	14	↓	3	19	152	21	228
Writtle	14*	↓	1	29	564	122	607
Silwood Park (nr Ascot)	47*	↓	/	5	139	/	61
Wye	51*	↓	/	19	348	/	92
Starcross (nr Exeter)	65*	↑	14	12	192	70	168

The mealy cabbage aphid was caught at ten sites this week, with hotspots at Wellesbourne (855), and Hereford (413). Accumulated numbers are above the 10-year averages at Wellesbourne, Kirton, Hereford and Wye.

Willow-carrot aphid (*Cavariella aegopodii*)

Willow-carrot aphid (<i>Cavariella aegopodii</i>)	Bulletin Week Totals 13/07-19/07				Accumulated until 19/07		
	2015	Compared to last Bulletin week	2014	10-year average 2005-14	2015	2014	10-year average 2005-14
Dundee	4*	↓	2	5	77	/	75
Gogarbank (Edinburgh)	3	↑	0	3	232	98	88
Newcastle	4*	↓	2	3	109	/	109
York	0*	↓	N/A	N/A	454	N/A	N/A
Preston	0*	↓	2	2	545	/	545
Kirton	8	↓	/	4	1978	/	612
Broom's Barn (nr Bury St Edmunds)	16	↑	0	7	1979	375	737
Wellesbourne	0*	↓	0	1	1169	/	434
Hereford	2*	↑	0	3	366	60	390
Rothamsted (Harpenden)	0		0	5	1412	103	380
Writtle	3*	↑	0	3	2400	148	833
Silwood Park (nr Ascot)	0*	↓	/	1	671	/	238
Wye	4*	↑	/	0	1073	/	360
Starcross (nr Exeter)	2*	↓	3	2	346	42	158

The willow-carrot aphid was caught at nine sites this week, with numbers highest at Broom's Barn (16). Numbers decreased at eight sites and are now low everywhere.

Pea aphid (*Acyrtosiphon pisum*)

Pea aphid (<i>Acyrtosiphon pisum</i>)	Bulletin Week Totals 13/07-19/07				Accumulated until 19/07		
	2015	Compared to last Bulletin week	2014	10-year average 2005-14	2015	2014	10-year average 2005-14
Dundee	2*	↓	12	16	6	/	58
Gogarbank (Edinburgh)	8	↑	6	4	17	26	38
Newcastle	0*		5	4	1	/	25
York	7*	↑	N/A	N/A	26	N/A	N/A
Preston	1*	↑	2	3	12	/	49
Kirton	18	↓	/	101	188	/	959
Broom's Barn (nr Bury St Edmunds)	14	↓	2	102	66	125	647
Wellesbourne	43*	↑	87	110	127	/	384
Hereford	12*	↑	14	34	52	37	141
Rothamsted (Harpenden)	24	↑	35	96	106	169	446
Writtle	25*	↑	20	117	148	264	786
Silwood Park (nr Ascot)	15*	↑	/	24	43	/	138
Wye	20*	↓	/	106	101	/	338
Starcross (nr Exeter)	25*	↑	9	19	260	86	133

The pea aphid was caught at thirteen sites this week, with numbers increasing slightly at nine sites. Accumulated numbers are below the 10yr averages everywhere except Starcross.

Further information

Please send information on crop aphids to: mark-s.taylor@rothamsted.ac.uk

AHDB Cereals and Oilseeds: [Click here](#)

AHDB Potatoes: [Click here](#)

AHDB Horticulture: [Click here](#)

Rothamsted Insect Survey: [Click here](#)

Science and Advice for Scottish Agriculture (SASA): [Click here](#)

In partnership with



© Agriculture and Horticulture Development Board 2015. All rights reserved.

While the Agriculture and Horticulture Development Board seeks to ensure that the information contained within this document is accurate at the time of printing, no warranty is given in respect thereof and, to the maximum extent permitted by law, the Agriculture and Horticulture Development Board accepts no liability for loss, damage or injury howsoever caused (including that caused by negligence) or suffered directly or indirectly in relation to information and opinions contained in or omitted from this document.

Reference herein to trade names and proprietary products without stating that they are protected does not imply that they may be regarded as unprotected and thus free for general use. No endorsement of named products is intended, nor is any criticism implied of other alternative, but unnamed, products.